

## Case Report

### **Penil Buschke–Lowenstein Tumor arising from common genital warts (a case report)**

**Darabi MR (MD)\*<sup>1</sup>, Khooei AR (MD)<sup>2</sup>, Kalani Moghaddam F (MD)<sup>3</sup>**

<sup>1</sup>*Professor, Department of Urology, Mashhad University of Medical Sciences, Mashhad, Iran.*

<sup>2</sup>*Associate Professor, Department of Pathology, Mashhad University of Medical Sciences, Mashhad, Iran.*

<sup>3</sup>*General Physician, Mashhad University of Medical Sciences, Mashhad, Iran.*

---

#### **Abstract**

Buschke–Lowenstein Tumor (BLT) or Giant Condyloma Acuminatum (GCA) is a slow-growing, locally aggressive, cauliflower-like tumor of great size that usually arises in the perineal region. Buschke–Lowenstein Tumor is triggered by human papilloma virus (HPV), usually either genotype 6 or 11. Buschke–Lowenstein Tumor is usually preceded by condyloma acuminatum and occurs at any age after puberty, usually between the 4<sup>th</sup> and 6<sup>th</sup> decades. Invasive overgrowth and recurrence after treatment are its characteristics and malignant transformation is also possible. Here, we have reported a case of a 33 year-old man with penile Buschke–Lowenstein Tumor arising from common genital warts.

**Keywords:** Buschke-Lowenstein Tumor, Genital Warts, Giant Condyloma Acuminatum, Penis

---

\* **Corresponding Author:** *Darabi MR (MD), E-mail: j\_darabi@yahoo.com*

**Received** 16 Sep 2008

**Revised** 23 Jan 2010

**Accepted** 9 Feb 2010